

Convergence of eigenvalues for a highly non-self-adjoint differential operator

E. B. Davies ^{*} John Weir [†]

September 4, 2008

Abstract

In this paper we study a family of operators dependent on a small parameter $\varepsilon > 0$, which arise in a problem in fluid mechanics. We show that the spectra of these operators converge to $\mathbf{N}$ as $\varepsilon \rightarrow 0$, even though, for fixed $\varepsilon > 0$, the eigenvalue asymptotics are quadratic.

1 Introduction

In a recent paper [1] Benilov, O'Brien and Sazonov argued that the equation

$$\frac{\partial f}{\partial t} = Hf \tag{1}$$

approximates the evolution of a liquid film inside a rotating horizontal cylinder, where H is the closure of the operator H_0 on $L^2(-\pi, \pi)$ defined by

$$(H_0 f)(\theta) = \varepsilon \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial f}{\partial \theta} \right) + \frac{\partial f}{\partial \theta} \tag{2}$$

for any sufficiently small fixed $\varepsilon > 0$ and all $f \in \text{Dom}(H_0) = \mathcal{C}_{\text{per}}^2([-\pi, \pi])$. They also made several conjectures, based on non-rigorous asymptotic and numerical analysis, including that the spectrum of H is purely imaginary and consists of eigenvalues which accumulate at $\pm i\infty$, and that the eigenvalues converge to $i\mathbf{Z}$ as $\varepsilon \rightarrow 0$. Weir proved these conjectures, except for the

^{*}Department of Mathematics, King's College London, Strand, London WC2R 2LS, e.brian.davies@kcl.ac.uk

[†]Department of Mathematics, King's College London, Strand, London WC2R 2LS, john.l.weir@kcl.ac.uk

convergence of the eigenvalues, in [2, 3]. We prove the remaining conjecture in this paper.

Davies showed in [4] that, for $0 < \varepsilon \leq 2$, every $\lambda \in \mathbf{C}$ is an eigenvalue of H . For $\varepsilon < 2$ he showed that $-iH$ has compact resolvent by considering the unitarily equivalent operator A on $l^2(\mathbf{Z})$ defined by

$$(Av)_n = \frac{\varepsilon}{2}n(n-1)v_{n-1} - \frac{\varepsilon}{2}n(n+1)v_{n+1} + nv_n \quad (3)$$

for all $v \in \text{Dom}(A) = \{v \in l^2(\mathbf{Z}) : Av \in l^2(\mathbf{Z})\}$. Here $A = \mathcal{F}^{-1}(-iH)\mathcal{F}$, where $\mathcal{F} : L^2(-\pi, \pi) \rightarrow l^2(\mathbf{Z})$ is the Fourier transform. If $\mathcal{F}f = v$ then $(v_n)_{n \in \mathbf{Z}}$ are the Fourier coefficients of f . This result was achieved by obtaining sharp bounds on the rate of decay of eigenvectors and resolvent kernels, and by determining the precise domains of the operators involved. He also showed that

$$A = A_- \oplus 0 \oplus A_+, \quad (4)$$

where A_- and A_+ are the restrictions of A to $l^2(\mathbf{Z}_-)$ and $l^2(\mathbf{Z}_+)$ respectively, and that A_- is unitarily equivalent to $-A_+$. Since the resolvent is compact and the adjoint has the same eigenvalues, the spectrum of $-iH$ consists entirely of eigenvalues.

Weir proved in [2] that these eigenvalues, if they exist, must all be real. Boulton, Levitin and Marletta subsequently proved in a recent paper [5] that a wider class of operators possess only real eigenvalues. However, they did not prove that any non-zero eigenvalues exist for these operators, nor that their spectra are real. In [3], Weir proved rigorously that $-iH$ has infinitely many eigenvalues, all of multiplicity one, which accumulate at $\pm\infty$ by showing that the eigenvalues of A_+ correspond to those of a self-adjoint operator with compact resolvent. This operator L_ε on $L^2((0, 1), 2w_\varepsilon(x)dx)$ is defined as the closure of the operator given by

$$(L_\varepsilon f)(x) = -\frac{\varepsilon}{2}w_\varepsilon(x)^{-1}(p_\varepsilon f')'(x) \quad (5)$$

for all $f \in \{f \in \mathcal{C}^\infty([0, 1]) : f(0) = 0\}$, where

$$\begin{aligned} w_\varepsilon(x) &= x^{-1}(1-x)^{1/\varepsilon}(1+x)^{-1/\varepsilon}, \\ p_\varepsilon(x) &= (1-x)^{1+1/\varepsilon}(1+x)^{1-1/\varepsilon}. \end{aligned}$$

It was argued in [6] that the distribution of the eigenvalues, if they exist, should be quadratic, but no rigorous bounds were given. By analysing the self-adjoint operator, Weir proved rigorously in [3] that $\lambda_n \sim \varepsilon\pi^2 n^2 \beta^{-2}$ as $n \rightarrow \infty$ for a certain explicit constant β .

The conjecture in [1] that $\lambda_{\varepsilon,n} \rightarrow n$ as $\varepsilon \rightarrow 0$ is supported by numerical evidence in the same paper, and also in [6, 4, 3]. In this paper we prove the conjecture rigorously. In Section 2 we apply unitary transformations to obtain a family of operators on the same space. These operators are invertible with Hilbert-Schmidt inverses (Theorem 2.2). We then identify a differential operator with spectrum $\mathbf{N}$ (Theorem 3.3, Corollary 3.4), which is unitarily equivalent to an operator whose inverse has an integral kernel which is the pointwise limit of the integral kernels of the aforementioned family of inverses (Theorem 3.6). Finally we show that we actually have norm convergence (Theorem 4.2) and use the variational method to show that this implies convergence of the eigenvalues, i.e. $\lambda_{\varepsilon,n} \rightarrow n$ as $\varepsilon \rightarrow 0$ for all $n \in \mathbf{N}$; see Theorem 4.3.

2 Unitary transformations

Since we are interested in the limit as $\varepsilon \rightarrow 0$, we assume for the rest of the paper that $0 < \varepsilon < 1$. In order to obtain convergence of operators in some sense, we need a family of operators on the same space. To this end we apply unitary transformations to L_ε to obtain a family of operators on $L^2((0, 1/\varepsilon), ds)$ and then extend each operator to an operator on $L^2((0, \infty), ds)$.

Lemma 2.1 *The operator L_ε , defined by (5), is unitarily equivalent to an operator $\tilde{L}_\varepsilon$ on $L^2((0, 1/\varepsilon), \tilde{w}_\varepsilon(s)ds)$ such that*

$$(\tilde{L}_\varepsilon g)(s) = -\tilde{w}_\varepsilon(s)^{-1}(\tilde{p}_\varepsilon g')(s) \quad (6)$$

where

$$\begin{aligned} \tilde{w}_\varepsilon(s) &= 2s^{-1}(1 - \varepsilon s)^{1/\varepsilon}(1 + \varepsilon s)^{-1/\varepsilon} \\ \tilde{p}_\varepsilon(s) &= (1 - \varepsilon s)^{1+1/\varepsilon}(1 + \varepsilon s)^{1-1/\varepsilon} \end{aligned}$$

for all $s \in (0, 1/\varepsilon)$. Moreover, $\tilde{L}_\varepsilon$ is invertible with inverse R_ε given by

$$(R_\varepsilon f)(s) = \int_0^{1/\varepsilon} G_\varepsilon(s, t) f(t) \tilde{w}_\varepsilon(t) dt \quad (7)$$

for all $f \in L^2((0, 1/\varepsilon), \tilde{w}_\varepsilon(s)ds)$, where

$$\gamma_\varepsilon(s) = \int_0^s \tilde{p}_\varepsilon(u)^{-1} du \quad (8)$$

$$= \frac{1}{2} \left\{ \left(\frac{1 + \varepsilon s}{1 - \varepsilon s} \right)^{1/\varepsilon} - 1 \right\} \quad (9)$$

and

$$G_\varepsilon(s, t) = \begin{cases} \gamma_\varepsilon(s) & \text{if } 0 < s \leq t < 1/\varepsilon \\ \gamma_\varepsilon(t) & \text{if } 0 < t \leq s < 1/\varepsilon. \end{cases} \quad (10)$$

Proof We define an operator $U : L^2((0, 1), 2w_\varepsilon(x)dx) \rightarrow L^2((0, 1/\varepsilon), \tilde{w}_\varepsilon(s)ds)$ by

$$(Uf)(s) = f(\varepsilon s) \quad (11)$$

for all $f \in L^2((0, 1), 2w_\varepsilon(x)dx)$, $s \in (0, 1/\varepsilon)$. It is easy to show that U is a unitary operator. We define $\tilde{L}_\varepsilon = UL_\varepsilon U^{-1}$. A simple calculation shows that (6) holds. The analogous result about the inverse of L_ε was proven in [3], so the result about $\tilde{L}_\varepsilon$ follows from the unitary equivalence of the two operators. It only remains for us to calculate

$$\begin{aligned} \gamma_\varepsilon(s) &= \int_0^s \tilde{p}_\varepsilon(u)^{-1} du \\ &= \frac{1}{\varepsilon} \int_0^{\varepsilon s} p_\varepsilon(u)^{-1} du \\ &= \frac{1}{\varepsilon} \int_0^{\varepsilon s} \frac{1}{1-u^2} \left(\frac{1+u}{1-u} \right)^{1/\varepsilon} du \\ &= \frac{1}{\varepsilon} \int_0^{\varepsilon s} \frac{1}{(1-u)^2} \left(\frac{1+u}{1-u} \right)^{1/\varepsilon-1} du \\ &= \frac{1}{2} \left[\left(\frac{1+u}{1-u} \right)^{1/\varepsilon} \right]_0^{\varepsilon s} \\ &= \frac{1}{2} \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s} \right)^{1/\varepsilon} - 1 \right\}. \end{aligned}$$

■

Theorem 2.2 *The operator L_ε is unitarily equivalent to an operator M_ε on $L^2((0, 1/\varepsilon), ds)$ such that*

$$(M_\varepsilon^{-1}f)(s) = \int_0^{1/\varepsilon} K_\varepsilon(s, t) f(t) dt \quad (12)$$

for all $f \in L^2((0, 1/\varepsilon), ds)$, where

$$K_\varepsilon(s, t) = \begin{cases} (st)^{-1/2} \left(\frac{1-\varepsilon s}{1+\varepsilon s} \right)^{1/2\varepsilon} \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s} \right)^{1/\varepsilon} - 1 \right\} \left(\frac{1-\varepsilon t}{1+\varepsilon t} \right)^{1/2\varepsilon} & \text{if } 0 \leq s \leq t \\ (st)^{-1/2} \left(\frac{1-\varepsilon s}{1+\varepsilon s} \right)^{1/2\varepsilon} \left\{ \left(\frac{1+\varepsilon t}{1-\varepsilon t} \right)^{1/\varepsilon} - 1 \right\} \left(\frac{1-\varepsilon t}{1+\varepsilon t} \right)^{1/2\varepsilon} & \text{if } 0 \leq t \leq s. \end{cases} \quad (13)$$

Moreover, if we define N_ε on $L^2((0, \infty), ds)$ by

$$(N_\varepsilon f)(s) = \int_0^\infty \tilde{K}_\varepsilon(s, t) f(t) dt \quad (14)$$

for all $f \in L^2((0, \infty), ds)$, where

$$\tilde{K}_\varepsilon(s, t) = \begin{cases} K_\varepsilon(s, t) & \text{if } 0 < s, t \leq 1/\varepsilon \\ 0 & \text{otherwise,} \end{cases} \quad (15)$$

then N_ε has the same non-zero eigenvalues as L_ε^{-1} , and each non-zero eigenvalue has the same multiplicity with respect to each operator.

Proof We define a unitary operator $J_\varepsilon : L^2((0, 1/\varepsilon), ds) \rightarrow L^2((0, 1/\varepsilon), \tilde{w}_\varepsilon(s) ds)$ by

$$(J_\varepsilon f)(s) = \tilde{w}_\varepsilon(s)^{-1/2} f(s) \quad (16)$$

and then put $M_\varepsilon = J_\varepsilon^{-1} \tilde{L}_\varepsilon J_\varepsilon$. We then have $M_\varepsilon^{-1} = J_\varepsilon^{-1} R_\varepsilon J_\varepsilon$, so

$$(M_\varepsilon^{-1} f)(s) = \int_0^{1/\varepsilon} K_\varepsilon(s, t) f(t) dt \quad (17)$$

for all $f \in L^2((0, 1/\varepsilon), ds)$, where

$$\begin{aligned} K_\varepsilon(s, t) &= \tilde{w}_\varepsilon(s)^{1/2} G_\varepsilon(s, t) \tilde{w}_\varepsilon(t)^{1/2} \\ &= \begin{cases} \tilde{w}_\varepsilon(s)^{1/2} \gamma_\varepsilon(s) \tilde{w}_\varepsilon(t)^{1/2} & \text{if } 0 < s \leq t < 1/\varepsilon, \\ \tilde{w}_\varepsilon(s)^{1/2} \gamma_\varepsilon(t) \tilde{w}_\varepsilon(t)^{1/2} & \text{if } 0 < t \leq s < 1/\varepsilon. \end{cases} \end{aligned}$$

Substituting in expression (9) for γ_ε , we obtain (13).

There is no unitary equivalence between M_ε^{-1} and N_ε , but it is easy to see that they have the same eigenvalues, and that the non-zero eigenvalues have the same multiplicities with respect to each operator. Indeed,

$$N_\varepsilon = M_\varepsilon^{-1} \oplus 0,$$

where 0 is the zero operator acting on $L^2((1/\varepsilon, \infty), ds)$. The theorem now follows from the unitary equivalence established in Lemma 2.1. ■

3 The limit operator

In this section we consider the operator L_0 on $\mathcal{H} = L^2((0, \infty), w_0(s) ds)$, where

$$w_0(s) = \lim_{\varepsilon \rightarrow 0} \tilde{w}_\varepsilon(s) = 2s^{-1} e^{-2s}, \quad (18)$$

defined on some suitable domain, which we identify below, by

$$(L_0 f)(s) = -w_0(s)^{-1}(p_0 f')(s), \quad (19)$$

where $p_0(s) = \lim_{\varepsilon \rightarrow 0} \tilde{p}_\varepsilon(s) = e^{-2s}$. We shall show that L_0 has a self-adjoint extension, which is invertible and whose inverse is Hilbert-Schmidt. We then identify the spectrum of $\tilde{L}_0$ and the integral kernel of its inverse.

Definition Let $\mathcal{P}$ denote the set of all polynomials on $(0, \infty)$ and $s\mathcal{P}$ denote those elements of $\mathcal{P}$ which have constant term zero.

Definition Let $\text{Dom}(L_0)$ be the set of twice differentiable functions $g : (0, \infty) \rightarrow \mathbf{C}$ such that $\lim_{s \rightarrow 0+} g(s) = 0$, $\limsup_{s \rightarrow 0+} |g'(s)| < \infty$, $\lim_{s \rightarrow \infty} e^{-s} g(s) = 0$, $\lim_{s \rightarrow \infty} e^{-s} g'(s) = 0$ and $L_0 f \in \mathcal{H}$.

Remark It is easy to show that $\text{Dom}(L_0) \subset \mathcal{H}$. We may therefore define L_0 on this domain by equation (19). Note also that $s\mathcal{P} \subset \text{Dom}(L_0)$.

Lemma 3.1 *The set $\mathcal{P}$ is dense in $L^2((0, \infty), 2se^{-2s}ds)$.*

Proof Let $g \in L^2((0, \infty), 2se^{-2s}ds)$ be such that $g \perp \mathcal{P}$ and define

$$f(s) = g(s)2se^{-2s}\chi_{(0, \infty)}(s)$$

for all $s \in \mathbf{R}$. We need to prove that $g = 0$, or equivalently that $f = 0$. We first prove that $f \in L^2(\mathbf{R})$ and the Fourier transform $\hat{f}$ of f has an analytic continuation to the set $\{z : |\Im z| < 1\}$. We then prove that $\hat{f}$ is zero on an interval containing zero, and hence is identically zero. The invertibility of the Fourier transform then implies that $f = 0$.

Firstly

$$\int_{\mathbf{R}} |f(s)|^2 ds \leq \sup_{s \in (0, \infty)} 2se^{-2s} \int_0^\infty |g(s)|^2 2se^{-2s} ds < \infty$$

so $f \in L^2(\mathbf{R})$. If $b < 1$ then a similar estimate shows that $e^{b|s|}f \in L^2(\mathbf{R})$. Theorem IX.13 of [7] implies that $\hat{f}$ has an analytic continuation to $\{z : |\Im z| < 1\}$.

In order to evaluate $\hat{f}$ in a neighbourhood of zero, we first approximate $e^{-i\xi x}$ by polynomials:

$$\left| e^{-i\xi x} - \sum_{n=0}^{N-1} \frac{(-i\xi x)^n}{n!} \right| \leq \sum_{n=N}^{\infty} \frac{|x|^n |\xi|^n}{n!}$$

$$\begin{aligned}
&= |x|^N |\xi|^N \sum_{n=0}^{\infty} \frac{|x|^n |\xi|^n}{(n+N)!} \\
&\leq \frac{|x|^N |\xi|^N}{N!} \sum_{n=0}^{\infty} \frac{|x|^n |\xi|^n}{n!} \\
&= \frac{|x|^N |\xi|^N}{N!} e^{|x\xi|}
\end{aligned}$$

for all $x \in \mathbf{R}$, $\xi \in \mathbf{R}$, $N \in \mathbf{N}$. We now choose $\beta \in (0, 1)$. If $\delta > 0$ is sufficiently small then $e^{(\beta+\delta)|x|} f \in L^2(\mathbf{R})$. Also $e^{-\delta|x|} \in L^2(\mathbf{R})$, so $e^{\beta|x|} f \in L^1(\mathbf{R})$. If $|\xi| < \beta/2$ then

$$\begin{aligned}
|\hat{f}(\xi)| &= \left| \int_{\mathbf{R}} e^{-i\xi x} f(x) dx \right| \\
&\leq \left| \int_{\mathbf{R}} \left\{ e^{-i\xi x} - \sum_{n=0}^{N-1} \frac{(-i\xi x)^n}{n!} \right\} f(x) dx \right| + \left| \int_{\mathbf{R}} \sum_{n=0}^{N-1} \frac{(-i\xi x)^n}{n!} f(x) dx \right| \\
&\leq \int_{\mathbf{R}} \frac{|x|^N |\xi|^N}{N!} e^{|x\xi|} |f(x)| dx + \left| \int_0^{\infty} \sum_{n=0}^{N-1} \frac{(-i\xi x)^n}{n!} g(x) 2x e^{-2x} dx \right| \\
&\leq \sup_{x \in (0, \infty)} \frac{|x|^N |\xi|^N}{N!} e^{(|\xi|-\beta)|x|} \int_{\mathbf{R}} |f(x)| e^{\beta|x|} dx
\end{aligned}$$

for all $N \in \mathbf{N}$. The final integral is finite since $e^{\beta|x|} f \in L^1(\mathbf{R})$. We put $h_{\xi, N}(x) = \frac{|x|^N |\xi|^N}{N!} e^{(|\xi|-\beta)|x|}$ for all $x \in (0, \infty)$ and note that this is a smooth, positive function of x . As $x \rightarrow 0$ or $x \rightarrow \infty$, $h(x) \rightarrow 0$, so the supremum of h must be obtained at a local maximum in the interval $(0, \infty)$. The only zero of h' is at $N/(\beta - |\xi|)$ so

$$\begin{aligned}
\sup_{x \in (0, \infty)} h(x) &= h(N/(\beta - |\xi|)) \\
&= \frac{N^N e^{-N}}{N!} \frac{|\xi|^N}{(\beta - |\xi|)^N} \\
&\sim (2\pi N)^{-1/2} \frac{|\xi|^N}{(\beta - |\xi|)^N} \\
&\rightarrow 0
\end{aligned}$$

as $N \rightarrow \infty$ for all $\xi \in [-\beta/2, \beta/2]$. Hence $\hat{f}(\xi) = 0$ for all $\xi \in [-\beta/2, \beta/2]$. ■

Corollary 3.2 *The set $s\mathcal{P}$ is dense in $\mathcal{H}$.*

Proof It is easy to show that the operator $U : L^2((0, \infty), 2se^{-s}ds) \rightarrow \mathcal{H}$ defined by

$$(Uf)(s) = sf(s)$$

for all $s \in (0, \infty)$ is unitary and that $U\mathcal{P} = s\mathcal{P}$. $\blacksquare$

Theorem 3.3 *The operator L_0 is symmetric and there is a complete orthonormal sequence $\{e_n\}_{n \in \mathbf{N}}$ in $\mathcal{H}$ such that, for each $n \in \mathbf{N}$, $e_n \in \text{Dom}(L_0)$ and $L_0 e_n = ne_n$.*

Proof For all $f, g \in \text{Dom}(L_0)$

$$\begin{aligned} \langle L_0 f, g \rangle &= - \int_0^\infty (p_0 f')'(s) \overline{g(s)} ds \\ &= p_0(0) f'(0) \overline{g(0)} - \lim_{n \rightarrow \infty} p_0(n) f'(n) \overline{g(n)} + \int_0^\infty p_0(s) f'(s) \overline{g'(s)} ds \\ &= \lim_{n \rightarrow \infty} p_0(n) f(n) \overline{g'(n)} - p_0(0) f(0) \overline{g'(0)} - \int_0^\infty f(s) \overline{(p_0 g')'(s)} ds \\ &= \langle f, L_0 g \rangle \end{aligned}$$

since $f(0) = g(0) = 0$,

$$\lim_{n \rightarrow \infty} p_0(n) f'(n) \overline{g(n)} = \left(\lim_{n \rightarrow \infty} e^{-n} f'(n) \right) \left(\lim_{n \rightarrow \infty} e^{-n} \overline{g(n)} \right) = 0$$

and similarly $\lim_{n \rightarrow \infty} p_0(n) f(n) \overline{g'(n)} = 0$. Therefore L_0 is symmetric.

For all $n \in \mathbf{N}$, define $a_{n,r}$ recursively for $r = 1, \dots, n$ by

$$a_{n,n} = 1 \tag{20}$$

$$a_{n,r} = -\frac{r(r+1)}{2(n-r)} a_{n,r+1} \text{ for } r = 1, \dots, n-1 \tag{21}$$

and put

$$f_n(s) = \sum_{r=1}^n a_{n,r} s^r \tag{22}$$

for all $s \in (0, \infty)$, $n \in \mathbf{N}$. Then, for each $n \in \mathbf{N}$, $f_n \in \text{Dom}(L_0)$ and

$$\begin{aligned} (L_0 f_n)(s) &= -\frac{p_0(s)}{w_0(s)} f_n''(s) - \frac{p_0'(s)}{w_0(s)} f_n'(s) \\ &= -\frac{s}{2} f_n''(s) + s f_n'(s) \\ &= -\sum_{r=2}^n \frac{r(r-1)a_{n,r}}{2} s^{r-1} + \sum_{r=1}^n r a_{n,r} s^r \\ &= \sum_{r=1}^{n-1} \left\{ r a_{n,r} - \frac{r(r+1)a_{n,r+1}}{2} \right\} s^r + n a_{n,n} s^n \\ &= n f_n(s) \end{aligned}$$

for all $s \in (0, \infty)$. Since the f_n are eigenvectors corresponding to distinct eigenvalues and L_0 is symmetric, they are orthogonal. Putting $e_n = f_n \|f_n\|^{-1}$, we obtain an orthonormal sequence such that $L_0 e_n = n e_n$. Each e_n is a polynomial of degree n , so $\text{lin}\{e_n\}_{n \in \mathbf{N}} = s\mathcal{P}$. It now follows from Corollary 3.2 that $\{e_n\}_{n \in \mathbf{N}}$ is complete. ■

The situation above can be seen as an exceptional case of that for the associated Laguerre polynomials $\{L_n^{(\alpha)}\}_{n=0}^{\infty}$ (see [8]). If $\alpha > -1$ these polynomials are a complete orthogonal set of eigenvectors, corresponding to eigenvalues $0, 1, 2, \dots$ respectively, of the differential equation

$$x f'' + (\alpha + 1 - x) f' + \lambda f = 0$$

on $L^2((0, \infty), x^\alpha e^{-x} dx)$. After a change of variables, the eigenvalue equation for L_0 becomes the exceptional case $\alpha = -1$. The singularity at the origin requires special treatment, but otherwise the treatment of this case is the same as for $\alpha > -1$.

Corollary 3.4 *The operator L_0 is essentially self-adjoint, and the spectrum of its closure is precisely $\mathbf{N}$.*

Proof The result follows immediately from the theorem and Lemma 1.2.2 of [9]. ■

Theorem 3.5 *The operator $\bar{L}_0$ is invertible and its inverse R is a Hilbert-Schmidt operator. Moreover,*

$$(R_0 f)(s) = \int_0^\infty G_0(s, t) f(t) w_0(t) dt \quad (23)$$

for all $f \in \mathcal{H}$, where

$$G_0(s, t) = \begin{cases} \gamma_0(s) & \text{if } 0 < s \leq t \\ \gamma_0(t) & \text{if } 0 < t \leq s \end{cases} \quad (24)$$

and

$$\begin{aligned} \gamma_0(s) &= \int_0^s p_0(u)^{-1} du \\ &= \int_0^s e^{2u} du \\ &= \frac{1}{2} \{e^{2s} - 1\}. \end{aligned}$$

Proof Let R_0 be defined by equation (23). Then R_0 is Hilbert-Schmidt since

$$\begin{aligned}
& \int_0^\infty \left(\int_0^\infty |G_0(s, t)|^2 w_0(t) dt \right) w_0(s) ds \\
&= 2 \int_0^\infty \left(\int_s^\infty |G_0(s, t)|^2 w_0(t) dt \right) w_0(s) ds \\
&= 2 \int_0^\infty \left(\int_s^\infty |e^{2s} - 1|^2 t^{-1} e^{-2t} dt \right) s^{-1} e^{-2s} ds \\
&\leq 2 \int_0^\infty \left(\int_s^\infty e^{-2t} dt \right) |e^{2s} - 1|^2 s^{-2} e^{-2s} ds \\
&= \int_0^\infty |e^{-2s} - 1|^2 s^{-2} ds \\
&\leq \int_0^1 4 ds + \int_1^\infty s^{-2} ds \\
&= 5.
\end{aligned}$$

Let $f \in s\mathcal{P}$. Then $f \in L^1((0, \infty), w_0(s) ds)$. Hence it is easy to show that Rf is absolutely continuous and

$$\begin{aligned}
(Rf)'(s) &= \int_s^\infty \gamma'(s) f(t) w_0(t) dt \\
&= p_0(s)^{-1} \int_s^\infty f(t) w_0(t) dt.
\end{aligned}$$

We now show that $Rf \in \text{Dom}(L_0)$. Since f , w_0 and p_0^{-1} are smooth on $(0, \infty)$, (Rf) is twice differentiable (indeed, it is also smooth). We calculate

$$\limsup_{s \rightarrow 0+} |(Rf)(s)| \leq \limsup_{s \rightarrow 0+} \gamma_0(s) \int_0^\infty |f(t)| w_0(t) dt = 0 \quad (25)$$

and

$$\begin{aligned}
\limsup_{s \rightarrow 0+} |(Rf)'(s)| &\leq \limsup_{s \rightarrow 0+} p_0(s)^{-1} \int_s^\infty |f(t)| w_0(t) dt \\
&= \int_0^\infty |f(t)| w_0(t) dt < \infty.
\end{aligned}$$

Since f is a polynomial, there exists a constant $c > 0$ such that $|f(s)| \leq ce^s$ for all $s \in (0, \infty)$. Let $\delta > 0$ be given. Then, for all $s > c/\delta$,

$$\begin{aligned}
|e^{-s}(Rf)(s)| &\leq e^{-s} \int_0^s |\gamma_0(t) f(t) w_0(t)| dt + e^{-s} \int_s^\infty |\gamma_0(s) f(t) w_0(t)| dt \\
&\leq e^{-s} \int_0^s e^{2t} |f(t)| t^{-1} e^{-2t} dt + e^{-s} \int_s^\infty e^{2s} |f(t)| t^{-1} e^{-2t} dt \\
&\leq e^{-s} \int_0^{c/\delta} |f(t)| t^{-1} dt + e^{-s} \int_{c/\delta}^s \delta e^t dt + s^{-1} e^s \int_s^\infty c e^{-t} dt \\
&\leq e^{-s} \int_0^{c/\delta} |f(t)| t^{-1} dt + \delta + c s^{-1} \rightarrow \delta
\end{aligned}$$

as $s \rightarrow \infty$. Since $\delta > 0$ is arbitrary, $e^{-s}(Rf)(s) \rightarrow 0$ as $s \rightarrow \infty$. Also

$$\begin{aligned} |e^{-s}(Rf)'(s)| &\leq e^s \int_s^\infty |f(t)| \frac{2}{t} e^{-2t} dt \\ &\leq 2s^{-1} e^s \int_s^\infty c e^{-t} dt \\ &= 2cs^{-1} \rightarrow 0 \end{aligned}$$

as $s \rightarrow \infty$. Therefore $Rf \in \text{Dom}(L_0)$, provided $L_0 Rf \in \mathcal{H}$, and

$$\begin{aligned} (\bar{L}_0 Rf)(s) &= (L_0 Rf)(s) \\ &= -w_0(s)^{-1} \frac{d}{ds} \int_s^\infty f(t) w_0(t) dt \\ &= f(s). \end{aligned}$$

Since $f \in \mathcal{H}$, we indeed have $L_0 Rf \in \mathcal{H}$.

We have proven that $\bar{L}_0 Rf = f$ for all $f \in s\mathcal{P}$. Let $f \in \mathcal{H}$. By Corollary 3.2, there is a sequence (f_n) in $s\mathcal{P}$ such that $f_n \rightarrow f$ as $n \rightarrow \infty$. By the above, we have $Rf_n \rightarrow Rf$ and $\bar{L}_0 Rf_n = f_n \rightarrow f$ as $n \rightarrow \infty$. Since $\bar{L}_0$ is closed, this implies that $Rf \in \text{Dom}(\bar{L}_0)$ and $\bar{L}_0 Rf = f$. Finally, $\bar{L}_0$ is injective since $0 \notin \text{Spec}(\bar{L}_0)$. Hence $\bar{L}_0 R\bar{L}_0 f = \bar{L}_0 f$ implies $R\bar{L}_0 f = f$ for all $f \in \mathcal{H}$. ■

Theorem 3.6 *The operator $\bar{L}_0$ is unitarily equivalent to an operator M_0 on $L^2((0, \infty), ds)$. Moreover,*

$$(M_0^{-1}f)(s) = \int_0^\infty K_0(s, t) f(t) dt \quad (26)$$

for all $f \in L^2((0, \infty), ds)$, where

$$\begin{aligned} K_0(s, t) &= w_0(s)^{1/2} G_0(s, t) w_0(t)^{1/2} \\ &= \begin{cases} (st)^{-1/2} e^{-s} \{e^{2s} - 1\} e^{-t} & \text{if } 0 < s \leq t \\ (st)^{-1/2} e^{-s} \{e^{2t} - 1\} e^{-t} & \text{if } 0 < t \leq s. \end{cases} \end{aligned}$$

Proof We define a unitary operator

$$J_0 : L^2((0, \infty), ds) \rightarrow L^2((0, \infty), w_0(s) ds)$$

by

$$(J_0 f)(s) = w_0(s)^{-1/2} f(s) \quad (27)$$

and put $M_0 = J_0^{-1} \bar{L}_0 J_0$. Then $M_0^{-1} = J_0^{-1} R_0 J_0$ and (26) is immediate. ■

4 Convergence of the eigenvalues

Clearly $\tilde{K}_\varepsilon(s, t) \rightarrow K_0(s, t)$ as $\varepsilon \rightarrow 0$ for each $s, t \in (0, \infty)$. In this section we shall show that $K_\varepsilon \rightarrow K_0$ in L^2 -norm as $\varepsilon \rightarrow 0$, and hence that $N_\varepsilon \rightarrow M_0^{-1}$ in Hilbert-Schmidt norm as $\varepsilon \rightarrow 0$. We use this fact to prove that the eigenvalues of L_ε converge to those of L_0 as $\varepsilon \rightarrow 0$.

Lemma 4.1 *If $0 \leq s \leq t < 1/\varepsilon$ then*

$$\left(\frac{1-\varepsilon s}{1+\varepsilon s}\right)^{1/2\varepsilon} \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s}\right)^{1/\varepsilon} - 1 \right\} \left(\frac{1-\varepsilon t}{1+\varepsilon t}\right)^{1/2\varepsilon} - e^{-s} \{e^{2s} - 1\} e^{-t} \leq e^{-s-t}. \quad (28)$$

Proof We first note that

$$\begin{aligned} \log \left(\left(\frac{1+\varepsilon s}{1-\varepsilon s}\right)^{1/2\varepsilon} \left(\frac{1-\varepsilon t}{1+\varepsilon t}\right)^{1/2\varepsilon} \right) &= \frac{1}{2\varepsilon} \{ \log(1+\varepsilon s) - \log(1-\varepsilon s) \} \\ &\quad + \frac{1}{2\varepsilon} \{ \log(1-\varepsilon t) - \log(1+\varepsilon t) \} \\ &= s - t + \sum_{k=1}^{\infty} \frac{\varepsilon^{2k}}{2k+1} (s^{2k+1} - t^{2k+1}) \\ &\leq s - t \end{aligned}$$

and hence

$$\left(\frac{1+\varepsilon s}{1-\varepsilon s}\right)^{1/2\varepsilon} \left(\frac{1-\varepsilon t}{1+\varepsilon t}\right)^{1/2\varepsilon} \leq e^{s-t} \quad (29)$$

for $0 \leq s \leq t < 1/\varepsilon$. For such s, t ,

$$\begin{aligned} \left(\frac{1-\varepsilon s}{1+\varepsilon s}\right)^{1/2\varepsilon} \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s}\right)^{1/\varepsilon} - 1 \right\} \left(\frac{1-\varepsilon t}{1+\varepsilon t}\right)^{1/2\varepsilon} &\leq \left(\frac{1+\varepsilon s}{1-\varepsilon s}\right)^{1/2\varepsilon} \left(\frac{1-\varepsilon t}{1+\varepsilon t}\right)^{1/2\varepsilon} \\ &\leq e^{s-t} \\ &= e^{-s} \{e^{2s} - 1\} e^{-t} + e^{-s-t}. \end{aligned}$$

■

Theorem 4.2 $\lim_{\varepsilon \rightarrow 0} \|N_\varepsilon - M_0^{-1}\|_{\text{HS}} = 0$.

Proof Using the symmetry of $\tilde{K}_\varepsilon$ and K_0 , it is sufficient to show that

$$\int_0^\infty \int_s^\infty |\tilde{K}_\varepsilon - K_0|^2 dt ds \rightarrow 0 \quad (30)$$

as $\varepsilon \rightarrow 0$. If $\frac{1}{2} \log 2 \leq s \leq t < 1/\varepsilon$ then, by Lemma 4.1,

$$\begin{aligned} st \left| \tilde{K}_\varepsilon - K_0 \right|^2 &= \left| \left(\frac{1-\varepsilon s}{1+\varepsilon s} \right)^{1/2\varepsilon} \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s} \right)^{1/\varepsilon} - 1 \right\} \left(\frac{1-\varepsilon t}{1+\varepsilon t} \right)^{1/2\varepsilon} - e^{-s} \{e^{2s} - 1\} e^{-t} \right|^2 \\ &\leq \max \left\{ e^{-2s-2t}, e^{-2s} \{e^{2s} - 1\}^2 e^{-2t} \right\} \\ &\leq e^{-2s} \{e^{2s} - 1\}^2 e^{-2t}. \end{aligned}$$

If $t \geq 1/\varepsilon$ then $\tilde{K}_\varepsilon = 0$, so the above bound still holds. Since

$$\begin{aligned} &\int_{\frac{1}{2} \log 2}^{\infty} \int_s^{\infty} (st)^{-1} e^{-2s} \{e^{2s} - 1\}^2 e^{-2t} dt ds \\ &\leq \int_{\frac{1}{2} \log 2}^{\infty} s^{-2} e^{-2s} \{e^{2s} - 1\}^2 \int_s^{\infty} e^{-2t} dt ds \\ &= \frac{1}{2} \int_{\frac{1}{2} \log 2}^{\infty} s^{-2} e^{-4s} \{e^{2s} - 1\}^2 ds \\ &\leq \frac{1}{2} \int_{\frac{1}{2} \log 2}^{\infty} s^{-2} ds \\ &< \infty, \end{aligned}$$

we may use the Lebesgue Dominated Convergence Theorem to prove that

$$\int_{\frac{1}{2} \log 2}^{\infty} \int_s^{\infty} \left| \tilde{K}_\varepsilon - K_0 \right|^2 dt ds \rightarrow 0 \quad (31)$$

as $\varepsilon \rightarrow 0$.

If $0 \leq s < 1$ and $0 < \varepsilon < 1$ then

$$\begin{aligned} \log \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s} \right)^{1/\varepsilon} \right\} &= 2 \sum_{k=0}^{\infty} \frac{\varepsilon^{2k} s^{2k+1}}{2k+1} \\ &\leq 2 \sum_{k=0}^{\infty} \frac{s^{2k+1}}{2k+1} \\ &= \log \left(\frac{1+s}{1-s} \right), \end{aligned}$$

so

$$\left(\frac{1+\varepsilon s}{1-\varepsilon s} \right)^{1/\varepsilon} \leq \log \left(\frac{1+s}{1-s} \right). \quad (32)$$

Also

$$\left(\frac{1-\varepsilon x}{1+\varepsilon x} \right)^{1/2\varepsilon} \leq e^{-x} \quad (33)$$

if $0 \leq x < 1/\varepsilon$. Hence, for $0 \leq s \leq \frac{1}{2} \log 2$, $s \leq t < 1/\varepsilon$,

$$\begin{aligned}
0 &\leq st \left| \tilde{K}_\varepsilon - K_0 \right|^2 \\
&\leq \max \left\{ \left(\frac{1-\varepsilon s}{1+\varepsilon s} \right)^{1/2\varepsilon} \left\{ \left(\frac{1+\varepsilon s}{1-\varepsilon s} \right)^{1/\varepsilon} - 1 \right\} \left(\frac{1-\varepsilon t}{1+\varepsilon t} \right)^{1/2\varepsilon}, e^{-s} \{e^{2s} - 1\} e^{-t} \right\}^2 \\
&\leq \max \left\{ e^{-2s} \left(\frac{1+s}{1-s} - 1 \right)^2 e^{-2t}, e^{-2s} \{e^{2s} - 1\}^2 e^{-2t} \right\}.
\end{aligned}$$

As before, this bound holds for all t such that $t \geq s$, since $\tilde{K}_\varepsilon = 0$ if $t \geq 1/\varepsilon$. Since we have

$$\begin{aligned}
&\int_0^{\frac{1}{2} \log 2} \int_s^\infty (st)^{-1} e^{-2s} \{e^{2s} - 1\}^2 e^{-2t} dt ds \\
&\leq \int_0^{\frac{1}{2} \log 2} s^{-2} e^{-2s} \{e^{2s} - 1\}^2 \int_s^\infty e^{-2t} dt ds \\
&= \frac{1}{2} \int_0^{\frac{1}{2} \log 2} s^{-2} e^{-4s} \{e^{2s} - 1\}^2 ds \\
&< \infty
\end{aligned}$$

and

$$\begin{aligned}
&\int_0^{\frac{1}{2} \log 2} \int_s^\infty (st)^{-1} e^{-2s} \left(\frac{1+s}{1-s} - 1 \right)^2 e^{-2t} dt ds \\
&= \int_0^{\frac{1}{2} \log 2} \int_s^\infty (st)^{-1} e^{-2s} \left(\frac{2s}{1-s} \right)^2 e^{-2t} dt ds \\
&\leq c \int_0^{\frac{1}{2} \log 2} \int_s^\infty e^{-2s} e^{-2t} dt ds \\
&= \frac{c}{2} \int_0^{\frac{1}{2} \log 2} e^{-4s} ds \\
&< \infty
\end{aligned}$$

for some constant c , we may use the Lebesgue Dominated Convergence Theorem to prove that

$$\int_0^{\frac{1}{2} \log 2} \int_s^\infty \left| \tilde{K}_\varepsilon - K_0 \right|^2 dt ds \rightarrow 0 \tag{34}$$

as $\varepsilon \rightarrow 0$. ■

We now use standard variational methods to deduce the convergence of the eigenvalues from the norm convergence of the resolvents.

Theorem 4.3 *For each $\varepsilon \in (0, 1)$, $n \in \mathbf{N}$, let $\lambda_{\varepsilon, n}$ be the n th eigenvalue of L_ε . Then $\lambda_{\varepsilon, n} \rightarrow n$ as $\varepsilon \rightarrow 0$.*

Proof If S is a finite-dimensional subspace of $L^2((0, \infty), ds)$ then we define

$$\mu_\varepsilon(S) = \sup \{ \langle (I - N_\varepsilon)f, f \rangle : f \in S \text{ and } \|f\| = 1 \}, \quad (35)$$

$$\mu_0(S) = \sup \{ \langle (I - S_0^{-1})f, f \rangle : f \in S \text{ and } \|f\| = 1 \}, \quad (36)$$

$$\mu_{\varepsilon,n} = \inf \{ \mu_\varepsilon(S) : S \subset L^2((0, \infty), ds) \text{ and } \dim(S) = n \}, \quad (37)$$

$$\mu_{0,n} = \inf \{ \mu_0(S) : S \subset L^2((0, \infty), ds) \text{ and } \dim(S) = n \}. \quad (38)$$

For each $\varepsilon \in (0, 1)$, N_ε is self-adjoint since it is a Hilbert-Schmidt operator with a symmetric integral kernel. Also $\text{Spec}(L_\varepsilon) \subset [1, \infty)$ and hence it follows from Theorem 2.2 that $I - N_\varepsilon$ is a non-negative self-adjoint operator with essential spectrum $\{1\}$ and eigenvalues $\{1 - 1/\lambda_{\varepsilon,n} : n \in \mathbf{N}\} \subset [0, 1)$. It follows from Corollary 3.4 and Theorem 3.6 that $I - M_0^{-1}$ is a non-negative self-adjoint operator with essential spectrum $\{1\}$ and eigenvalues $\{1 - 1/n : n \in \mathbf{N}\} \subset [0, 1)$.

By Theorem 4.5.2 of [9], $\mu_{\varepsilon,n} = 1 - 1/\lambda_{\varepsilon,n}$ and $\mu_{0,n} = 1 - 1/n$. Let $\delta > 0$ be given. Then Theorem 4.2 implies that there exists $\eta > 0$ such that, whenever $0 < \varepsilon < \eta$,

$$\langle (I - M_0^{-1})f, f \rangle - \delta \leq \langle (I - N_\varepsilon)f, f \rangle \leq \langle (I - M_0^{-1})f, f \rangle + \delta \quad (39)$$

for all $f \in L^2((0, \infty), ds)$ such that $\|f\| = 1$. For such ε , this implies that

$$\mu_\varepsilon(S) - \delta \leq \mu_0(S) \leq \mu_\varepsilon(S) + \delta \quad (40)$$

for all finite-dimensional subspaces S of $L^2((0, \infty), ds)$ and hence that

$$\mu_{\varepsilon,n} - \delta \leq \mu_{0,n} \leq \mu_{\varepsilon,n} + \delta \quad (41)$$

for all $n \in \mathbf{N}$. Therefore, for all $n \in \mathbf{N}$, $\mu_{\varepsilon,n} \rightarrow \mu_{0,n}$ as $\varepsilon \rightarrow 0$ and hence $\lambda_{\varepsilon,n} \rightarrow n$ as $\varepsilon \rightarrow 0$. ■

References

- [1] E. S. Benilov, S. B. G. O'Brien, and I. A. Sazonov. A new type of instability: explosive disturbances in a liquid film inside a rotating horizontal cylinder. *Journal of Fluid Mechanics*, 497 (2003) 201–224.
- [2] J. L. Weir. An indefinite convection-diffusion operator with real spectrum. *Applied Mathematics Letters*, in press.
- [3] J. L. Weir. Correspondence of the eigenvalues of a non-self-adjoint operator to those of a self-adjoint operator. *Preprint*, 2008.

- [4] E. B. Davies. An indefinite convection-diffusion operator. *LMS Journal of Computation and Mathematics*, 10 (2007) 288–306.
- [5] Lyonell Boulton, Michael Levitin, and Marco Marletta. A PT-symmetric periodic problem with boundary and interior singularities. *arXiv:0801.0172*, 2008.
- [6] M. Chugunova and D. Pelinovsky. Spectrum of a non-self-adjoint operator associated with the periodic heat equation. *arXiv:math-ph/0702100*, 2007.
- [7] M. Reed and B. Simon. *Methods of Modern Mathematical Physics vol. II: Fourier Analysis, Self Adjointness*. Academic Press, 1975.
- [8] I. S. Gradshteyn and M. Ryzhik. *Tables of Integral Series and Products*. Academic Press, 1980.
- [9] E. B. Davies. *Spectral Theory and Differential Operators*. Cambridge University Press, 1995.